

Client	HEALTH BENCHMARKS, INC. STANDARD ALGORITHM <i>Implemented for Blue Cross Blue Shield of Texas</i>		
Measure Title	USE OF LONG-TERM CONTROL DRUGS FOR PERSISTENT ASTHMA		
Disease State	Asthma	Indicator Classification	Disease Management
Strength of Recommendation	A (<i>inhaled corticosteroid or inhaled corticosteroid combos</i>) B (<i>other classes of medication [i.e., mast cell stabilizers, leukotriene modifiers, methylxanthines]</i>)		
Organizations Providing Recommendation	Joint Council of Allergy, Asthma and Immunology National Asthma Education and Prevention Program National Heart, Lung and Blood Institute		
Clinical Intent	To ensure that members with persistent asthma receive medication appropriate for long term control of asthma.		
Physician Specialties	Family Practice, Geriatric Medicine, Internal Medicine, Pediatric Pulmonology, Pediatrics, Pulmonary Diseases		
Background	<p>Disease Burden</p> <ul style="list-style-type: none"> • An estimated 32.6 million persons in the United States have been diagnosed with asthma sometime in their lifetime, as of 2005.[1] • In 2004, asthma led to over 1.8 million emergency department visits, nearly 500,000 hospitalizations, and over 4,000 deaths.[1] <p>Reason for Indicated Intervention or Treatment</p> <ul style="list-style-type: none"> • Regular use of inhaled corticosteroids improves asthma control, and decreases hospital admissions and mortality from asthma in adults and children with persistent asthma.[2-4] • For patients with moderate persistent asthma, adding a long-acting beta-2 agonist to a low or medium dose inhaled corticosteroid improves lung function and symptoms, decreases asthma exacerbations, and reduces the use of additional short-acting beta-2-agonists.[4] • Many patients with persistent asthma are still being undertreated with long-term control medications.[5-7] <p>Evidence Supporting Intervention or Treatment</p> <ul style="list-style-type: none"> • Randomized, controlled trials have shown that inhaled corticosteroid use in patients with persistent asthma, when compared to placebo or beta-2-agonists, results in improved pre-bronchodilator FEV1, reduced oral steroid and supplemental short-acting beta-2-agonist use, and decreased airway responsiveness, asthma symptom scores, and hospitalizations.[8-23] 		

- Results from randomized, controlled trials on using leukotriene modifiers alone for those with persistent asthma are mixed. Some randomized control trials show no difference between leukotriene modifier and inhaled corticosteroid use [24-28], but others found increased asthma exacerbations and poorer symptom control in those using the leukotriene modifiers.[29-31]
- For asthma that is poorly controlled with inhaled corticosteroid use alone, randomized controlled trials have shown that patients have better symptom control when long-acting beta-2-agonists are added, instead of leukotriene modifiers.[32-34]
- Most randomized control trials demonstrate that adding a long-acting beta-2-agonist to an inhaled corticosteroid decreases asthma exacerbations more than increasing the inhaled corticosteroid dose.[35-38] However, one randomized control trial found that increasing the inhaled corticosteroid dose led to better symptom control than adding a long-acting beta-2-agonist.[39]

Clinical Recommendations

- An expert panel convened by the National Heart, Lung and Blood Institute and the National Asthma Education and Prevention Program (NAEPP) developed Expert Panel Report 3 (EPR3) Guidelines for the Diagnosis and Management of Asthma published in 2007. EPR3 states that the following medications are appropriate for the long-term control of persistent asthma:[40]
 - Corticosteroids – most potent and effective medication currently available
 - Cromolyn sodium and nedocromil – used as alternative but not preferred for treatment of persistent asthma.
 - Immunomodulators – Omalizumab (anti-IgE) as adjunctive therapy for severe persistent asthma[4]
 - Leukotriene modifiers – alternative, but not preferred therapy for the treatment of mild asthma
 - Long acting beta agonist – should not be used as monotherapy – but can be used in combination with inhaled corticosteroids for long-term control and prevention of systems in moderate or severe persistent asthma
 - Methylxathines – used as alternative, not preferred, adjunctive therapy with inhaled corticosteroid

Source Healthcare Effectiveness Data and Information Set (HEDIS®) 2008 Technical Specification for Physician Measurement

Denominator Definition Continuously enrolled members ages 5 - 56 years with evidence of persistent asthma (mild to severe) who meet at least 1 of the following criteria in *both* the measurement year and the year prior to the measurement year (criteria need not be the same across both years):

- At least 4 medication dispensing events

- Members whose 4 asthma medication dispensing events were solely for leukotriene modifiers must have a diagnosis of asthma during the same year as the leukotriene modifier (i.e. the measurement year or the year prior) or additionally meet any of the other persistent asthmatic criteria to be considered in the denominator
 - At least 1 ED visit with a primary diagnosis of asthma
 - At least 1 acute inpatient discharge with asthma as the primary diagnosis
 - At least 4 outpatient visits with asthma as one the listed diagnoses and at least two medication dispensing events

Denominator Codes Asthma
 ICD-9 diagnosis code(s): 493.xx
ED setting
 CPT-4 code(s): 99281-99285
 UB revenue code(s): 045x, 0981
Acute inpatient encounter
 CPT-4 code(s): 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 99261-99263, 99291
 UB revenue code(s): 010x, 0110-0114, 0119, 0120-0124, 0129, 0130-0134, 0139, 0140-0144, 0149, 0150-0154, 0159, 016x, 020x-022x, 072x, 0987
Outpatient encounter
 CPT-4 code(s): 99201-99205, 99211-99215, 99217-99220, 99241-99245, 99341-99345, 99347-99350, 99382-99386, 99392-99396, 99401-99404, 99411, 99412, 99420, 99429, 99499
 UB revenue code(s): 051x, 0520-0523, 0526-0529, 057x-059x, 077x, 0982, 0983
Acute inpatient encounter
 CPT-4 code(s): 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 99261-99263, 99291
 UB revenue code(s): 010x, 0110-0114, 0119, 0120-0124, 0129, 0130-0134, 0139, 0140-0144, 0149, 0150-0154, 0159, 016x, 020x-022x, 072x, 0987

Denominator Exclusion Definition Members who were diagnosed with emphysema or chronic obstructive pulmonary disease (COPD) any time prior to the end of the measurement year.

Denominator Exclusion Codes Emphysema
 ICD-9 diagnosis code(s): 492.x, 506.4, 518.1, 518.2
COPD
 ICD-9 diagnosis code(s): 491.2x, 493.2, 496, 506.4

Numerator Definition Members who received a prescription for a medication appropriate for long-term control of asthma during the measurement year.

Numerator Codes N/A

Physician Attribution Description **If client data does not contain PCP:**

Score all physicians (in the selected specialties) who saw the member during the measurement year.

If client data does contain PCP:

Score all primary care physicians who were assigned to the member during the measurement year.

References

1. Akinbami, L.J., *Asthma Prevalence, Health Care Use and Mortality: United States, 2003-05*, in *NCHS Health and Stats*. 2007, National Center for Health and Statistics: Hyattsville, MD.
2. Donahue, J.G., et al., *Inhaled steroids and the risk of hospitalization for asthma*. *Jama*, 1997. **277**(11): p. 887-91.
3. Suissa, S., et al., *Low-dose inhaled corticosteroids and the prevention of death from asthma*. *N Engl J Med*, 2000. **343**(5): p. 332-6.
4. *National Asthma Education and Prevention Program Expert Panel Report: guidelines for the diagnosis and management of asthma update on selected topics-2002*. *J Allergy Clin Immunol* 2002. **110**(5 pt 2): p. S141-219.
5. Jatulis, D.E., et al., *Preventive pharmacologic therapy among asthmatics: five years after publication of guidelines*. *Ann Allergy Asthma Immunol*, 1998. **81**(1): p. 82-8.
6. Wu, A.W., et al., *Quality of care and outcomes of adults with asthma treated by specialists and generalists in managed care*. *Arch Intern Med*, 2001. **161**(21): p. 2554-60.
7. *Use of Appropriate Medications for People with Asthma*. , in *The State of Managed Care Quality*. 2004, National Committee for Quality Assurance.
8. *Long-term effects of budesonide or nedocromil in children with asthma. The Childhood Asthma Management Program Research Group*. *N Engl J Med*, 2000. **343**(15): p. 1054-63.
9. Hoekstra, M.O., et al., *Eosinophil and mast cell parameters in children with stable moderate asthma*. *Pediatr Allergy Immunol*, 1998. **9**(3): p. 143-9.
10. Jonasson, G., K. Carlsen, and P. Blomqvist, *Clinical efficacy of low-dose inhaled budesonide once or twice daily in children with mild asthma not previously treated with steroids*. . *Eur Respir J* 1998. **12**(5): p. 1099-104.
11. Simons, F.E., *A comparison of beclomethasone, salmeterol, and placebo in children with asthma. Canadian Beclomethasone Dipropionate-Salmeterol Xinafoate Study Group*. *N Engl J Med*, 1997. **337**(23): p. 1659-65.
12. van Essen-Zandvliet, E.E., et al., *Effects of 22 months of treatment with inhaled corticosteroids and/or beta-2-agonists on lung function, airway responsiveness, and symptoms in children with asthma. The Dutch*

- Chronic Non-specific Lung Disease Study Group*. *Am Rev Respir Dis*, 1992. **146**(3): p. 547-54.
13. Adams, N., J. Bestall, and P.W. Jones, *Budesonide for chronic asthma in children and adults*. *Cochrane Database Syst Rev*, 2001(4): p. CD003274.
 14. Adams, N., J. Bestall, and P.W. Jones, *Inhaled fluticasone propionate for chronic asthma*. *Cochrane Database Syst Rev*, 2001(3): p. CD003135.
 15. Adams, N.P., et al., *Inhaled beclomethasone versus placebo for chronic asthma*. *Cochrane Database Syst Rev*, 2005(1): p. CD002738.
 16. Banov, C.H., W.C. Howland, 3rd, and W.R. Lumry, *Once-daily budesonide via Turbuhaler improves symptoms in adults with persistent asthma*. *Ann Allergy Asthma Immunol*, 2001. **86**(6): p. 627-32.
 17. Bronsky, E., et al., *Comparative clinical study of inhaled beclomethasone dipropionate and triamcinolone acetonide in persistent asthma*. *Ann Allergy Asthma Immunol*, 1998. **80**(4): p. 295-302.
 18. Kemp, J., et al., *Rapid onset of control with budesonide Turbuhaler in patients with mild-to-moderate asthma*. *Ann Allergy Asthma Immunol*, 1999. **82**(5): p. 463-71.
 19. McFadden, E.R., et al., *Administration of budesonide once daily by means of turbuhaler to subjects with stable asthma*. *J Allergy Clin Immunol*, 1999. **104**(1): p. 46-52.
 20. Miyamoto, T., et al., *A double-blind, placebo-controlled dose-response study with budesonide Turbuhaler in Japanese asthma patients. Japanese Pulmicort Turbuhaler study group*. *Respirology*, 2000. **5**(3): p. 247-56.
 21. O'Byrne, P.M., et al., *Low dose inhaled budesonide and formoterol in mild persistent asthma: the OPTIMA randomized trial*. *Am J Respir Crit Care Med*, 2001. **164**(8 Pt 1): p. 1392-7.
 22. Pauwels, R.A., et al., *Early intervention with budesonide in mild persistent asthma: a randomised, double-blind trial*. *Lancet*, 2003. **361**(9363): p. 1071-6.
 23. Wolfe, J.D., et al., *Effectiveness of fluticasone propionate in patients with moderate asthma: a dose-ranging study*. *Clin Ther*, 1996. **18**(4): p. 635-46.
 24. Busse, W., et al., *Low-dose fluticasone propionate compared with montelukast for first-line treatment of persistent asthma: a randomized clinical trial*. *J Allergy Clin Immunol*, 2001. **107**(3): p. 461-8.
 25. Israel, E., et al., *Effects of montelukast and beclomethasone on airway function and asthma control*. *J Allergy Clin Immunol*, 2002. **110**(6): p. 847-54.
 26. Nathan, R.A., E.R. Bleeker, and C. Kalberg, *A comparison of short-term treatment with inhaled fluticasone propionate and zafirlukast for patients with persistent asthma*. *Am J Med*, 2001. **111**(3): p. 195-202.
 27. Riccioni, G., et al., *Effectiveness of montelukast versus budesonide on quality of life and bronchial reactivity in subjects with mild-persistent asthma*. *Int J Immunopathol Pharmacol*, 2002. **15**(2): p. 149-155.
 28. Riccioni, G., et al., *Comparison of montelukast and budesonide on bronchial reactivity in subjects with mild-moderate persistent asthma*. *Pulm Pharmacol Ther*, 2003. **16**(2): p. 111-4.

29. Baumgartner, R.A., et al., *Distribution of therapeutic response in asthma control between oral montelukast and inhaled beclomethasone*. Eur Respir J, 2003. **21**(1): p. 123-8.
30. Brabson, J.H., et al., *Efficacy and safety of low-dose fluticasone propionate compared with zafirlukast in patients with persistent asthma*. Am J Med, 2002. **113**(1): p. 15-21.
31. Meltzer, E.O., et al., *Efficacy and safety of low-dose fluticasone propionate compared with montelukast for maintenance treatment of persistent asthma*. Mayo Clin Proc, 2002. **77**(5): p. 437-45.
32. Chervinsky, P., et al., *Comparison of mometasone furoate administered by metered dose inhaler with beclomethasone dipropionate*. Int J Clin Pract, 2002. **56**(6): p. 419-25.
33. Fish, J.E., et al., *Salmeterol powder provides significantly better benefit than montelukast in asthmatic patients receiving concomitant inhaled corticosteroid therapy*. Chest, 2001. **120**(2): p. 423-30.
34. Ringdal, N., et al., *The salmeterol/fluticasone combination is more effective than fluticasone plus oral montelukast in asthma*. Respir Med, 2003. **97**(3): p. 234-41.
35. Baraniuk, J., et al., *Fluticasone alone or in combination with salmeterol vs triamcinolone in asthma*. Chest, 1999. **116**(3): p. 625-32.
36. Price, D., et al., *Early asthma control and maintenance with eformoterol following reduction of inhaled corticosteroid dose*. Thorax, 2002. **57**(9): p. 791-8.
37. Shrewsbury, S., S. Pyke, and M. Britton, *Meta-analysis of increased dose of inhaled steroid or addition of salmeterol in symptomatic asthma (MIASMA)*. Bmj, 2000. **320**(7246): p. 1368-73.
38. Woolcock, A., et al., *Comparison of addition of salmeterol to inhaled steroids with doubling of the dose of inhaled steroids*. Am J Respir Crit Care Med, 1996. **153**(5): p. 1481-8.
39. Pauwels, R.A., et al., *Effect of inhaled formoterol and budesonide on exacerbations of asthma. Formoterol and Corticosteroids Establishing Therapy (FACET) International Study Group*. N Engl J Med, 1997. **337**(20): p. 1405-11.
40. NAEPP. *Expert Panel Report 3: Guidelines for the Diagnosis and Management of Asthma, Full Report 2007*. 2007 2008 14 January [cited; Available from: <http://www.nhlbi.nih.gov/guidelines/asthma/asthgdln.pdf>.
41. *Global Strategy for Asthma Management and Prevention* Global Initiative for Asthma. p. 110.