





## Adult Health - for ages 19 and over

Preventive care is important. Making good, basic health choices can boost your health and well being. Some of these positive choices include:

- Eating a healthy diet
- Not using tobacco
- Striving for a healthy weight

- Getting regular exercise
- Limiting alcohol use

Learn more. For additional preventive care information, visit:

- ahrq.gov/patients-consumers/prevention/index.html
- cancer.org/cancer/risk-prevention/diet-physical-activity.html
- cdc.gov/vaccines/

Blue Cross and Blue Shield of Texas, a Division of Health Care Service Corporation, a Mutual Legal Reserve Company, an Independent Licensee of the Blue Cross and Blue Shield Association

752510.1223

## **Preventive Care**

Men and women are encouraged to get care as needed, follow a healthy lifestyle and make regular screenings a priority for better overall health. Discuss recommended preventive care services, screenings and immunizations with your health care provider. Don't hesitate to ask about nutrition and exercise, losing weight and stopping smoking. Other topics to discuss may include:

- Dental health
- Problems with drugs or alcohol
- Sexual behavior and sexually transmitted infections
- Feelings of depression
- Domestic violence
- Accident/injury prevention
- Preventing falls

Screenings		
Weight	Every visit or at least annually	
Body Mass Index (BMI)	Every visit or at least annually	
Blood Pressure (BP)	Every visit or at least annually	
Cholesterol	Adults 40 to 75 years of age should be screened; or adults 20 to 39 years old who have risk for coronary heart disease. Talk with your health care provider <sup>*</sup> about the starting and frequency of screening that is best for you.	
Colon Cancer Screening	Adults age 45-75 for colorectal cancer using: Guaiac Fecal Occult Blood Test (gFOBT) annually or; Fecal Immunochemical Testing (FIT) annually or; Fecal Immunochemical Testing (FIT)-DNA every 1-3 years or; Flexible sigmoidoscopy every 5 years or; Flexible sigmoidoscopy every 10 years with FIT annually or; Colonoscopy every 10 years or; CT Colonography every 5 years. <sup>#</sup> The risks and benefits of different screening methods vary. For details about pharmacy benefit coverage, call the number on the back of your member ID card.	
Diabetes Screening	Those with high blood pressure should be screened. Those who are overweight or have cardiovascular risk factors should be screened. All others should be screened starting at age 45. <sup>#</sup>	
Hepatitis C (HCV) Screening	Once for adults age 19-79. Most adults need to be screened only once. Persons with continued risk for HCV infection (eg, PWID) should be screened periodically; and persons at high risk for infection	
HIV Screening	Adults ages 19 to 65, older adults at increased risk and all pregnant women should be screened. Those 26-45 years of age, should discuss their options with their health care provider.	
*A health care provider could be a doctor, primary care provider, physician assistant, nurse practitioner or		

other health care professional.

<sup>#</sup>Recommendations may vary. Discuss the start and frequency of screenings with your health care provider, especially if you are at increased risk.

Immunizations (Vaccines)		
Tetanus Diphtheria Pertussis (Td/Tdap)	Get Tdap vaccine once, then a Td booster every 10 years	
Influenza (Flu)	Yearly	
Human Papillomavirus (HPV)	All Adults age 19-26, 2 or 3 doses depending on age at time of initial vaccination if not already given. <sup>#</sup> Those 27-45 should discuss options with their health care provider.	
Herpes Zoster (Shingles)	Two doses of RZV starting at age 50, or one dose of ZVL at age 60 or over. Discuss your options with your health care provider.*	
Hepatitis B (Hep B)	2, 3, or 4 doses depending on vaccine or condition beginning at age 19-59. Discuss your options with your health care provider.*	
Varicella (Chicken Pox)	Two doses for adults born in 1980 or later who have not previously received the varicella vaccine or have no evidence of immunity.	
Pneumococcal (Pneumonia)	Ages 65 and over, one dose of PCV 15 followed by PPSV 23 OR one dose of PCV 20. Discuss your options with your health care provider.*	
Measles, Mumps, Rubella (MMR)	1 or 2 doses for adults born in 1957 or later who have no evidence of immunity	
COVID-19 Vaccine	The CDC recommends adults get the COVID-19 vaccine. Talk to your health care provider or pharmacist about the COVID-19 vaccine and when you should get it.	
*Recommendations may vary. Discuss screening options with your health care provider, especially if you are at increased risk.		

\*Coverage for preventive care services at no cost share may vary depending on your specific benefit plan and use of network providers. For questions, please call the Customer Service number on the back of your ID card.

Women's Recommendations		
Mammogram	At least every 2 years for women ages 50 to 74 Ages 40 to 49 should discuss the risks and benefits of screening with their health care provider	
Cervical Cancer Screening	Women ages 21 to 65: Pap test every 3 years Another option for ages 30 to 65: Pap test with HPV test every 5 years Women who have had a hysterectomy or are over age 65 may not need a Pap test <sup>*</sup>	
Osteoporosis Screening	Women who are at an increased risk for osteoporosis should be screened at ages 65 and older. Bone measurement testing is recommended for postmenopausal women younger than 65 years who are at increased risk of osteoporosis as determined by a formal clinical risk assessment tool.	
Intensive Behavioral Counseling	All sexually active individuals (12 years old and above) who are at increased risk for sexually transmitted infections (STIs).	

\*Recommendations may vary. Discuss screening options with your health care provider, especially if you are at increased risk.

Men's Recommendations		
Prostate Cancer Screening	Discuss the benefits and risks of screening with your health care provider. <sup>#</sup>	
Abdominal Aortic Aneurysm	Have an ultrasound once between ages 65 to 75 if you have ever smoked.	
Intensive Behavioral Counseling	All sexually active individuals (12 years old and above) who are at increased risk for sexually transmitted infections (STIs).	
#Coverage for preventive care services at no cost share may vary depending on your specific benefit plan and use of		

<sup>#</sup>Coverage for preventive care services at no cost share may vary depending on your specific benefit plan and use of network providers. For questions, please call the Customer Service number on the back of your ID card.

The recommendations provided in the table are based on information from organizations such as the Advisory Committee on Immunization Practices, the American Academy of Family Physicians, the American Cancer Society and the United States Preventive Services Task Force. The recommendations are not intended as medical advice nor meant to be a substitute for the individual medical judgment of a health care provider. Please check with your health care provider for individualized advice on the recommendations provided.

## To ask for auxiliary aids and services or materials in other formats and languages at no cost, please call 1-866-689-1523 (TTY/TDD: 711).

Blue Cross and Blue Shield of Texas complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex.

ATENCIÓN: Si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1-855-710-6984 (TTY: 711).

CHÚ Ý: Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 1-855-710-6984 (TTY: 711).